

Tristan Juhan, MD

Orthopedic Surgeon

Office: 408-412-8100

Email: tjuhan@goldenstateortho.com

Website: tristanjuhanMD.com



**BICEPS TENDON REPAIR
REHABILITATION PROTOCOL**

	RANGE OF MOTION	IMMOBILIZER	THERAPEUTIC EXERCISE
PHASE I 0 - 3 weeks	None	Locked in neutral - worn at all times	Gentle wrist and shoulder ROM
PHASE II 3 - 6 weeks	30 degrees to 60 degrees in brace, very gradual progression of extension	Worn at all times (including exercise) - removed for hygiene	Continue with wrist and shoulder ROM, begin active extension to 30°, NO active flexion, gentle joint mobilizations
PHASE III 6 - 9 weeks	Active extension to 0 in brace	Worn at all times (including exercise) - removed for hygiene	Continue to maintain wrist and shoulder flexibility, begin rotator cuff/deltoid isometrics, progress active extension in brace
PHASE IV 9 - 12 weeks	Gently advance ROM to tolerance	None	Begin active flexion and extension against gravity, advance strengthening in phase III to resistive, maintain flexibility/ROM
PHASE V 12 weeks - 6 months	Gradual return to full and pain-free	None	Begin gentle flexion strengthening, advance activities in phase IV
PHASE VI 6 months and beyond	Full and pain-free	None	Return to full activity